

LIA™

Calibrachoa hybrida



LIA™ Spark Lavender



- Selected for basket production
- Blooms in a 10 hour day
- **Tested for pH sensitivity**
- Mature height and width
12 inches x 14 inches
(30cm x 36cm)



LIA™ Chili Pepper

The LIA™ series:



LIA™ Abstract
Lemon Cherry



LIA™ Sky Blue



LIA™ Violet



LIA™ Spark Pink



LIA™ Blue



LIA™ Bubblegum



LIA™ Dark Red



LIA™ Glaze



LIA™ Melon



LIA™ Abstract Pink



LIA™ Cranberry



LIA™ Yellow Imp.

EU & ROW
ONLY

For additional technical growing information:

LIA™ [CLICK HERE](#)



 **DANZIGER**

www.danzigeronline.com |   

LIA™

Calibrachoa hybrida



PROPAGATION TIPS

- Stick on priority – Number 3 out of 4 categories



Flowering time



Width



Height

Spring / Summer
Flowering

(12 - 14")
30 - 36 cm

(12 - 14")
30 - 36 cm

Average Time	Temperature	Hormone	Fertilization	Fungicide
5 weeks	Weeks 1 - 2 70°-72° F (21° - 22°C)	Optional	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 5 65° - 68° (18° - 20° C)		Weeks 3-5 100 to 150 ppm N	Day of sticking & Week 2



Irrigation specification - Remove from mist as soon as possible



Application of K-IBA can hasten rooting. Light in propagation for optimal growth and to hasten crop time. Daminozide at 1200 to 1500 ppm day of sticking. Daminozide plus Uniconazole-P tank mix week 3 to control stretch if needed.

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.2 / 5.8	0.80	72 / 74 F (22 / 23 C)	50 ppm N in mist			Day of sticking	Feeding through Mist
Week 2	5.2 to 5.8	.80 to .90	72 / 74 F (22 / 23 C)	50 ppm N in mist			Second fungicide app	
Week 3	5.2 to 5.8	.90 to 1.0	Cool to 68 / 70 F (20 / 21 C)	Feed 100 ppm to 150 ppm		Daminozide as needed		
Week 4	5.2 to 5.8	1.0 to 1.2	65 to 68 F (18 to 20 C)	Feed 100 ppm to 150 ppm		Daminozide as needed		
Week 5	5.2 to 5.8	1.0 to 1.2	65 to 68 F (18 to 20 C)	Feed 100 ppm to 150 ppm	Optional	Daminozide as needed	Week 5	

For additional technical growing information:

LIA™ [CLICK HERE](#)



DANZIGER
www.danzigeronline.com |



FINISHING TIPS

- Pinch is optional
- Cool temperature finishing for best quality

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
7 to 8 wks (12 cm) 5" pot	Average Day 68° to 72°F (20° to 22°C)	No pinch necessary	100 - 150 ppm N	In liner and finish stages
8 to 9 wks (15 cm) 6 ½"	Weeks 3 – 5 65° - 68° (18° - 20° C)	No lighting necessary	Soil EC 1.0 - 1.2 pH 5.5 to 6.0	
8 to 9 wks (20 cm) 8"				
8 to 9 wks (25 cm) 10"	8 to 9 wks 3 plants per container			

Pests	Diseases
Aphids	Botrytis
Thrip	Pythium
	Phytophthora
ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS	CYPRODINIL+FLUDIOXONIL, IPRDIONE, POLYOXIN
METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD	PROPAMOCARB
	PROPAMOCARB

- ! • Additional Iron (Fe) or Manganese (Mn) may be required depending on water quality and alkalinity
- Scout for Thrip on regular basis

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 6 Transplant	5.6 to 6.0	1.0 to 1.2	65 to 68 F (18 to 20 C)	feed 100 to 150 ppm	Day Neutral	Daminozide as needed	Spray fungicide after transplant	
Week 6 to 13	5.6 to 6.0	1.0 to 1.2	65 to 68 F (18 to 20 C)	feed 100 to 150 ppm	Not Required	Paclobutrazol Drench at 1 to 2 ppm		Paclobutrazol 1 to 2 ppm drench when plants reach 85% of desired size

For additional technical growing information:

LIA™ [CLICK HERE](#)



LIA™

Calibrachoa hybrida



Sawaya Waterford 2022



For additional technical growing information:

LIA™ [CLICK HERE](#)



 **DANZIGER**
www.danzigeronline.com |   